

Restore Foam Dressing without Border with TRIACT ADVANCED Technology



Protocol

This protocol is intended to supplement, not replace, your facility's guidelines and/or common practices.

Desired Outcomes

- To maintain a moist wound environment for the healing of partial to full-thickness wounds
- To help prevent pain during dressing removal
- To absorb exudate in moderate to heavy exuding wounds

Equipment

1. Restore Foam Dressing without Border
2. Wound measuring guide
3. Restore Wound Cleanser or sterile saline for wound cleansing
4. Gauze or towel to dry periwound skin
5. Secondary cover dressing
6. Gloves

Procedures

- Preparation of wound area
- Application of wound dressing
- Removal of wound dressing
- Documentation of observations

Preparation of Wound Area

1. Assemble supplies and take to bedside.
2. Identify and inform patient of procedure.
3. Provide for privacy.
4. Wash hands and apply gloves.
5. Remove dressing if present.
6. Thoroughly rinse or irrigate the wound area with Restore Wound Cleanser or sterile saline, if appropriate. *(If necessary, the wound should be debrided).*
7. Clean and dry the periwound skin.
8. Measure the wound using the wound measuring guide.
9. Choose a dressing size which ensures the pad will cover the entire wound.

Application of Dressing

1. Remove clear plastic tabs from the dressing.
2. Apply Restore Foam Dressing without Border (micro-adherent side down) to cover the entire wound using aseptic technique. The dressing may overlap onto healthy skin.
3. Secure with appropriate secondary dressing (gauze, transparent film, etc) using aseptic technique.
4. Remove gloves and wash hands after completing procedure.

Removal of Dressing

1. Wash hands and put on gloves.
2. Remove secondary dressing.
3. Remove Restore Foam Dressing without Border.
4. Irrigate wound base using Restore Wound Cleanser or sterile saline.
5. Reapply dressing if necessary.
6. Remove gloves and wash hands after completing procedure.

Note: Restore Foam Dressing without Border may be left in place for up to seven days. Dressing change frequency will depend on patient condition and the level of exudate.

Documentation of Observations

Document dressing changes, wound measurements and other pertinent observations regarding the wound or periwound site.

Assess

Key observations to document with wound dressing change:

- Size of wound
- Depth of tissue involvement (partial-thickness, full-thickness, stage)
- Presence of undermining or tunnelling
- Anatomic location
- Wound Base (granulation, epithelialization, subcutaneous tissue, muscle, bone or other underlying structures, non-viable/necrotic tissue, color, exudate amount/odor/color)
- Edge of wound
- Presence of foreign bodies
- Condition of periwound skin

Plan

Select interventions to provide appropriate local wound care:

- Prevent and manage infection
- Pain management
- Cleanse wound
- Remove non-viable tissue
- Manage exudate
- Eliminate dead space
- Control odor
- Protect wound and periwound skin
- Maintain a moist wound environment

Implement

Follow Protocol for the Restore Foam Dressing without Border

- Reduce or eliminate factors that may impede wound healing
- Provide systemic support for wound healing (proper nutrition, etc.)

Evaluate

- Reassess wound with each dressing change.
- Frequency of dressing changes should be dictated by facility protocol.
- If no progress can be demonstrated within two to four weeks, reevaluate the overall treatment plan, as well as adherence to this plan, making modifications as necessary.

See Instructions for Use for important information regarding the use of this product at www.hollisterwoundcare.com/products/ifus.html

Contraindications

Restore Foam Dressing without Border should not be used on individuals who are sensitive to or who have had an allergic reaction to the dressing or one of its components.

Hollisterwoundcare and wave logo are trademarks of Hollister Incorporated. Restore, TRIACT ADVANCED, the TRIACT ADVANCED graphic are trademarks of Hollister Wound Care LLC. Made and printed in USA.
©2010 Hollister Wound Care LLC.

Restore Foam Dressing without Border Advantages

- Maintenance of a moist wound environment
- Minimizes or eliminates trauma to the wound bed upon dressing removal
- Helps reduce or eliminate pain at dressing changes
- Super absorbent foam pad provides vertical absorption of wound exudate and helps protect the surrounding skin from maceration
- Suitable for use under compression bandages
- Provides gentle micro-adherence to the periwound skin for one-handed application

Ordering Information

Restore Foam Dressing without Border	Stock No	Unit of Sale
--------------------------------------	----------	--------------

2.5" x 2.5" (6 cm x 6 cm)	509380	10
4" x 4" (10 cm x 10 cm)	509381	10
6" x 6" (15 cm x 15 cm)	509382	10
Heel 4.7" x 7.5" (12 cm x 19 cm)	509384	10

Restore Wound Cleanser

8 oz. (236 mL) Trigger Spray Bottle, Non-Sterile	529975	12/box
12 oz. (354 mL) Trigger Spray Bottle, Non-Sterile	529976	12/box



Manufactured for
Hollister Wound Care LLC
1580 South Milwaukee Avenue
Suite 405
Libertyville, Illinois 60048
1.888.740.8999

Distributed in Canada by
Hollister Limited
95 Mary Street
Aurora, Ontario L4G 1G3
1.800.263.7400

www.hollisterwoundcare.com